arXiv:1209.2528v1 [math.CO] 12 Sep 2012

On the spectral moment of graphs with k cut edges*

Shuchao Li[a,†], Huihui Zhang[a], Minjie Zhang[b]

[a]Faculty of Mathematics and Statistics, Central China Normal University, Wuhan 430079, P.R. China

[b]School of Mathematics and Physics, Hubei Institute of Technology, Huangshi 435003, P.R. China

Abstract: Let $A(G)$ be the adjacency matrix of a graph G with $\lambda_1(G)$, $\lambda_2(G)$, $\ldots$, $\lambda_n(G)$ being its eigenvalues in non-increasing order. Call the number $S_k(G) := \sum_{i=1}^{n} \lambda_i^k(G)\,(k = 0, 1, \ldots, n-1)$ the kth spectral moment of G. Let $S(G) = (S_0(G), S_1(G), \ldots, S_{n-1}(G))$ be the sequence of spectral moments of G. For two graphs G_1 and G_2, we have $G_1 \prec_s G_2$ if $S_i(G_1) = S_i(G_2)\,(i = 0, 1, \ldots, k-1)$ and $S_k(G_1) < S_k(G_2)$ for some $k \in \{1, 2, \ldots, n-1\}$. Denote by $\mathscr{G}_n^k$ the set of connected n-vertex graphs with k cut edges. In this paper, we determine the first, the second, the last and the second last graphs, in an S-order, among $\mathscr{G}_n^k$, respectively.

Keywords: Spectral moment; Cut edge; Clique

AMS subject classification: 05C50, 15A18

1. Introduction

All graphs considered here are finite, simple and connected. Undefined terminology and notation may be referred to [1]. Let $G = (V_G, E_G)$ be a simple undirected graph with n vertices. $G - v$, $G - uv$ denote the graph obtained from G by deleting vertex $v \in V_G$, or edge $uv \in E_G$, respectively (this notation is naturally extended if more than one vertex or edge is deleted). Similarly, $G + uv$ is obtained from G by adding an edge $uv \notin E_G$. For $v \in V_G$, let $N_G(v)$ (or $N(v)$ for short) denote the set of all the adjacent vertices of v in G and $d_G(v) = |N_G(v)|$, and $\mathrm{dist}_G(u, v)$ is the distance between u and v. For an edge subset E' of G, denoted by $G[E']$ the subgraph induced by E'. A cut edge in a connected graph G is an edge whose deletion breaks the graph into two components. Let $\mathscr{G}_n^k$ be the set of all n-vertex graphs, each of which contains k cut edges.

Let $A(G)$ be the adjacency matrix of a graph G with $\lambda_1(G), \lambda_2(G), \ldots, \lambda_n(G)$ being its eigenvalues in non-increasing order. The number $\sum_{i=1}^{n} \lambda_i^k(G)\,(k = 0, 1, \ldots, n-1)$ is called the kth spectral moment of G, denoted by $S_k(G)$. Let $S(G) = (S_0(G), S_1(G), \ldots, S_{n-1}(G))$ be the sequence of spectral moments of G. For two graphs G_1, G_2, we shall write $G_1 =_s G_2$ if $S_i(G_1) = S_i(G_2)$ for $i = 0, 1, \ldots, n-1$. Similarly, we have $G_1 \prec_s G_2$ (G_1 comes before G_2 in an S-order) if for some $k\,(1 \leq k \leq n-1)$, we have $S_i(G_1) = S_i(G_2)\,(i = 0, 1, \ldots, k-1)$ and $S_k(G_1) < S_k(G_2)$. We shall also write $G_1 \preceq_s G_2$ if $G_1 \prec_s G_2$ or $G_1 =_s G_2$. S-order has been used in producing graph catalogs (see [5]), and for a more general setting of spectral moments one may be referred to [4].

Investigation on S-order of graphs attracts more and more researchers' attention. Cvetković and Rowlinson [6] studied the S-order of trees and unicyclic graphs and characterized the first and the last graphs, in an S-order, of all trees and all unicyclic graph with given girth, respectively. Chen, Liu and Liu [2] studied the lexicographic ordering by spectral moments (S-order) of unicyclic graph with a given girth. Wu and Fan [12] determined the first and the last graphs, in an S-order, of all unicyclic graphs and bicyclic graphs, respectively. Pan et al. [11] gave the first $\sum_{k=1}^{\lfloor \frac{n-1}{3} \rfloor} (\lfloor \frac{n-k-1}{2} \rfloor - k + 1)$ graphs apart from an n-vertex path, in an S-order, of all trees with n vertices. Wu and Liu [13] determined the last $\lfloor \frac{d}{2} \rfloor + 1$, in

---

*Financially supported by the National Natural Science Foundation of China (Grant Nos. 11071096, 11271149) and the Special Fund for Basic Scientific Research of Central Colleges (CCNU11A02015).

†E-mail: lscmath@mail.ccnu.edu.cn (S.C. Li), 425333559@qq.com (H. Zhang)

an S-order, among all n-vertex trees of diameter $d\,(4 \leq d \leq n-3)$. Pan et al. [10] identified the last and the second last graphs, in an S-order, of quasi-trees. Hu, Li and Zhang [8] studied the spectral moments of graphs with given clique number and chromatic number, respectively. Li and Song [9] identified the last n-vertex tree with a given degree sequence in an S-order. Consequently, the last trees in an S-order in the sets of all trees of order n with the largest degree, the leaves number, the independence number and the matching number was also determined, respectively.

In light of the information available from the related results on the spectral moments of graphs, it is natural to consider this problem on some other class of graphs, and the connected graphs with k cut edges are a reasonable starting point for such a investigation. The n-vertex connected graphs with k cut edges have been considered in different fields [7, 14, 15, 16], whereas to our best knowledge, the spectral moments of graphs in $\mathscr{G}_n^k$ were, so far, not considered. Here, we identified the first, the second, the last and the second last graphs, in an S-order, among $\mathscr{G}_n^k$, respectively.

Throughout the text we denote by $P_n, K_{1,n-1}, C_n$ and K_n the path, star, cycle and complete graph on n vertices, respectively. Let $K_{1,n-1}^*$ be a graph obtained from a star $K_{1,n-1}$ by attaching a leaf to one leaf of $K_{1,n-1}$, U_n be a graph obtained from C_{n-1} by attaching a leaf to one vertex of C_{n-1}, and B_4, B_5 be two graphs obtained from two cycle C_3, C_3' of length 3 by identifying one edge of C_3 with one edge of C_3' and identifying one vertex of C_3 with one vertex of C_3', respectively; see Fig. 1.


B_4 B_5 K_6^1 K_6^2 K_6^3

P_6^1 P_6^2 P_6^3 $K(a_0, \{a_1, a_2, \ldots, a_k\})$

Figure 1: Graphs $B_4, B_5, K_6^1, K_6^2, K_6^3, P_6^1, P_6^2, P_6^3$ and $K(a_0, \{a_1, a_2, \ldots, a_k\})$.

The graph K_n^k is an n-vertex graph obtained by attaching k pendant vertices to one vertex of K_{n-k}. The graph P_n^k is a graph obtained by identifying one end-vertex of P_{k+1} with one vertex of C_{n-k}. For example, for $n = 6, K_6^0 = K_6, K_6^5$ is a star, $P_6^0 = C_6$ and $K_6^1, K_6^2, K_6^3, P_6^1, P_6^2, P_6^3$ are depicted in Fig. 1. In general, $K_n^0 = K_n, K_n^{n-1}$ is star $K_{1,n-1}$, $K_n^{n-2} \cong K_n^{n-1}$ and $P_n^0 = C_n$. Let $K(a_0, \{a_1, a_2, \ldots, a_k\})$ be a graph obtained from $K_{1,k}$ by replacing each $u_i \in V_{K_{1,k}}$ by a clique K_{a_i} $(a_i \geq 1, i = 0, 1, 2, \ldots, k)$; see Fig. 1. Denote

$$\mathscr{K}_n^k = \left\{ K(a_0, \{a_1, a_2, \ldots, a_k\}) : a_i \geq 1 (0 \leq i \leq k), \sum_{i=0}^{k} a_i = n \right\}.$$

Let F be a graph. An F-subgraph of G is a subgraph of G which is isomorphic to the graph F. Let $\phi_G(F)$ (or $\phi(F)$) be the number of all F-subgraph of G.

Lemma 1.1. (see [3]) *The kth spectral moment of G is equal to the number of closed walks of length k.*

Lemma 1.2. *For every graph G, we have*

(i) $S_4(G) = 2\phi(P_2) + 4\phi(P_3) + 8\phi(C_4)$ (see [5]);

(ii) $S_5(G) = 30\phi(C_3) + 10\phi(U_4) + 10\phi(C_5)$ (see[13]);

(iii) $S_6(G) = 2\phi(P_2)+12\phi(P_3)+6\phi(P_4)+12\phi(K_{1,3})+12\phi(U_5)+36\phi(B_4)+24\phi(B_5)+24\phi(C_3)+48\phi(C_4)+12\phi(C_6)$ (see[2]).

Lemma 1.3 ([3]). *Given a connected graph G, $S_0(G) = n, S_1(G) = l, S_2(G) = 2m, S_3(G) = 6t$, where n, l, m, t denote the number of vertices, the number of loops, the number of edges and the number of triangles contained in G, respectively.*

Lemma 1.4 ([6]). *In an S-order of the n-vertex unicyclic graphs with girth g, the first graph is U_n^g which is obtained by the coalescence of a cycle C_g with a path P_{n-g+1} at one of its end-vertices.*

2. The last and the second last graphs in an S-order among $\mathscr{G}_n^k$

In this section, we will determine the last two graphs, in an S-order, among $\mathscr{G}_n^k$. Let $\mathbb{E} = \{e_1, e_2, \ldots, e_k\}$ be the set of the cut edges of $G \in \mathscr{G}_n^k$. Note that $S_2(G) = 2|E_G|$, hence $S_2(G + e) > S_2(G)$. By Lemma 1.3, in order to determine the last graph in an S-order among $\mathscr{G}_n^k$, it suffices to choose graph $G \in \mathscr{G}_n^k$ such that its $S_2(G)$ is as large as possible. So we can have the following assumption throughout this section.

Assumption 0. Each component of $G - \mathbb{E}$ is a clique.

Theorem 2.1. *Of all the connected graphs with n vertices and k cut edges, the last graph in an S-order is obtained uniquely at K_n^k.*

Proof. If $k = 0$, then by Assumption 0 we have $\mathscr{G}_n^0 = \{K_n\}$, our result holds immediately. Therefore we may assume that $k \geq 1$. Again by Assumption 0, we can denote the components of $G - \mathbb{E}$ by $K_{a_0}, K_{a_1}, \ldots, K_{a_k}$, $a_0 + a_1 + \cdots + a_k = n$. Assume, without loss of generality, that $a_0 \geq a_1 \geq a_2 \geq \cdots \geq a_k \geq 1$.

Let $V_i = \{v \in V_{K_{a_i}} : v$ is an end-vertex of a cut edge of $G\}$. Choose $G \in \mathscr{G}_n^k$ such that G is as large as possible under the order $\preceq_s$. In order to complete the proof, it suffices to show the following facts.

Fact 1. $|V_i| = 1$ for $i = 0, 1, 2, \ldots, k$.

Proof. Suppose to the contrary that there exists $i \in \{0, 1, 2, \ldots, k\}$ such that $|V_i| > 1$. Let $u, u' \in V_{a_i}$, both u and u' are end-vertices of the cut edges of G. Denote $N_G(u) \backslash N_{K_{a_i}}(u) = \{w_1, w_2, \ldots, w_s\}$ and $N_G(u') \backslash N_{K_{a_i}}(u') = \{z_1, z_2, \ldots, z_l\}$. It is routine to check that $s \geq 1$, $l \geq 1$. Let

$$G^* = G - \{u'z_1, u'z_2, \ldots, u'z_l\} + \{uz_1, uz_2, \ldots, uz_l\},$$

then $G^* \in \mathscr{G}_n^k$.

On the one hand, $S_i(G) = S_i(G^*)$ for $i = 0, 1, 2, 3$. On the other hand, $\phi_G(P_2) = \phi_{G^*}(P_2), \phi_G(C_4) = \phi_{G^*}(C_4)$, hence by Lemma 1.2(i),

$$S_4(G) - S_4(G^*) = 4(\phi_G(P_3) - \phi_{G^*}(P_3)) = 4\left(\binom{s}{2} + \binom{l}{2} - \binom{s+l}{2}\right) = -4sl < 0.$$

which implies that $G \prec_s G^*$, a contradiction. Therefore $|V_i| = 1$ for $0 \leq i \leq k$. □

By Fact 1, we can assume that $V_i = \{u_i\}$ for $i = 0, 1, 2, \ldots, k$.

Fact 2. $G \in \mathscr{K}_n^k$.

Proof. If not, then there exists a cut edge $u_0u_i \in \mathbb{E}$ such that u_i is an end-vertex of another cut edge(s). Let

$$|N_G(u_i) \setminus (N_{K_{a_i}}(u_i) \cup \{u_0\})| = l, \quad |N_G(u_0) \setminus (N_{K_{a_0}}(u_0) \cup \{u_i\})| = s.$$

It is straightforward to check that $l \geq 1$ and $s \geq 0$.

First consider that $s \geq 1$. In this case, let

$$G^* = G - \{u_iz : z \in N_G(u_i) \setminus (N_{K_{a_i}}(u_i) \cup \{u_0\})\} + \{u_0z : z \in N_G(u_i) \setminus (N_{K_{a_i}}(u_i) \cup \{u_0\})\}.$$

It is easy to see that $G^* \in \mathscr{G}_n^k$. Note that $S_i(G) = S_i(G^*)$ for $i = 0, 1, 2, 3$ and $\phi_G(P_2) = \phi_{G^*}(P_2)$, $\phi_G(C_4) = \phi_{G^*}(C_4)$, hence

$$S_4(G) - S_4(G^*) = 4(\phi_G(P_3) - \phi_{G^*}(P_3)) = 4(l(a_i - 1) - l(a_0 - 1) - ls) = 4l(a_i - a_0 - s) < 0,$$

which implies that $G \prec_s G^*$, a contradiction.

Now consider that $s = 0$. In this case, there exists a cut edge $u_iu_j \in \mathbb{E}$ such that u_j is an end-vertex of another cut edge(s). Let $|N_G(u_j) \setminus (N_{K_{a_j}}(u_j) \cup \{u_i\})| = p$. It is straightforward to check that $p \geq 1$. Let

$$\begin{aligned} G^* &= G - \{u_jz : z \in N_G(u_j) \setminus (N_{K_{a_j}}(u_j) \cup \{u_i\})\} + \{u_0z : z \in N_G(u_j) \setminus (N_{K_{a_j}}(u_j) \cup \{u_i\})\} \\ &\quad - \{u_iw : w \in N_G(u_i) \setminus (N_{K_{a_i}}(u_i) \cup \{u_0\})\} + \{u_0w : w \in N_G(u_i) \setminus (N_{K_{a_i}}(u_i) \cup \{u_0\})\}. \end{aligned}$$

It is easy to see that $G^* \in \mathscr{G}_n^k$. Note that $S_i(G) = S_i(G^*)$, $i = 0, 1, 2, 3$ and $\phi_G(P_2) = \phi_{G^*}(P_2)$, $\phi_G(C_4) = \phi_{G^*}(C_4)$. Hence,

$$\begin{aligned} S_4(G) - S_4(G^*) &= 4(\phi_G(P_3) - \phi_{G^*}(P_3)) \\ &= 4l(a_i - 1) + p(a_j - 1) - 4p(l - 1) - 4p - 4(l + p)(a_0 - 1) \\ &= 4[l(a_i - a_0) + p(a_j - a_0) - pl] < 0. \end{aligned}$$

The last inequality follows from $a_i \leq a_0, a_j \leq a_0$ and $pl > 0$. Hence, we obtain that $G \prec_s G^*$, a contradiction. Therefore $G \in \mathscr{K}_n^k$. □

By Fact 2, we can assume that $u_0u_j \in \mathbb{E}$, $1 \leq j \leq k$.

Fact 3. $a_1 = a_2 = \cdots = a_k = 1$.

Proof. Assume to the contrary that there exists a $j \in \{1, 2, \ldots, k\}$ such that $a_j > 1$. By Fact 2, we have $G = K(a_0, \{a_1, \ldots, a_{j-1}, a_j, a_{j+1}, \ldots, a_k\})$. Now we consider $G^* = K(a_0 + a_j - 1, \{a_1, \ldots, a_{j-1}, 1, a_{j+1}, \ldots, a_k\})$. It is easy to see that $G^* \in \mathscr{K}_n^k$.

Note that $S_i(G) = S_i(G^*)$, $i = 0, 1$ and

$$S_2(G) - S_2(G^*) = 2(a_j - 1) - 2(a_j - 1)a_0 = 2(a_j - 1)(1 - a_0) < 0,$$

i.e., $G \prec_s G^*$, a contradiction. Therefore $a_j = 1$ for $j = 1, 2, \ldots, k$. □

In the view of Fact 3, we have $a_0 = n - k$. Hence, $G = K(n - k, \{1, 1, \ldots, 1\})$, i.e., $G \cong K_n^k$, as desired. □

In the rest of this section, we are to determine last graph in an S-order among $\mathscr{G}_n^k \backslash K_n^k$. Delete an edge, say xy, from K_n and denote the resultant graph by G_1. Let G_2 be a graph obtained from G_1 by attaching a pendant vertex to one vertex, say r, of G_1 with $r \neq x, y$. Let $G_3 = K(n-k, \{\underbrace{1, 1, \ldots, 1}_{k}\}) - uw + vw$, where uw is a cut edge and u, v are two different vertices in $V_{K_{n-k}}$.

Based on Lemma 1.3, it is easy to see that among $\mathscr{G}_n^0$, K_n (resp. G_1) is the last (resp. the second last) graph in an S-order, while among $\mathscr{G}_n^1$ with $n \geq 5$, based on $S_2(G)$, the second last graph in an S-order must be a graph obtained from K_n^1 by deleting a non-cut edge, say e, from K_n^1. Denote the resultant graph by G' if e has a common vertex with the cut edge in K_n^1 and by G_2 otherwise. Note that $S_i(G_2) = S_i(G')$ for $i = 0, 1, 2, 3$ and $\phi_{G_2}(P_2) = \phi_{G'}(P_2)$, $\phi_{G_2}(C_4) = \phi_{G'}(C_4)$, hence by Lemma 1.2(i)

$$S_4(G') - S_4(G_2) = 4(\phi_{G'}(P_3) - \phi_{G_2}(P_3)) = -4 < 0,$$

i.e., $G' \prec_s G_2$, Hence, among $\mathscr{G}_n^1$ with $n \geq 5$, G_2 is the second last graph in an S-order. In what follows we only consider $k \geq 2$.

Theorem 2.2. *Among $\mathscr{G}_n^k$ with $2 \leq k \leq n-1$, the second last graph in an S-order is obtained uniquely at G_3 if $k \in \{2, 3, \ldots, n-2\}$ and at $K_{1,n-1}^*$ otherwise, where G_3 is defined as above.*

Proof. Choose $G \in \mathscr{G}_n^k \setminus \{K_n^k\}$ such that it is as large as possible according to $\preceq_s$. Denote the components of $G - \mathbb{E}$ by $U_0, U_1, U_2, \ldots, U_k$. We are to show that each of the components is a complete graph. In fact, if there exists a U_i which is not a complete graph, i.e., U_i contains two vertices x, y satisfying $xy \notin E_{U_i}$. Let $G' = G + xy$. If $G' \ncong K_n^k$, it is easy to see that $G \prec_s G'$, a contradiction. If $G' \cong K_n^k$, then either x or y is not an end-vertex of a cut edge of G. Without loss of generality, assume that x is not an end-vertex of a cut edge of G, delete a cut edge of G' and connect the isolated vertex with x by an edge; denote the resultant graph by G''. Then we have $S_0(G) = S_0(G''), S_1(G) = S_1(G'')$ and $S_2(G) < S_2(G'')$. Hence, $G \prec_s G''$, a contradiction. Therefore, we may denote the components of $G - \mathbb{E}$ by $K_{a_0}, K_{a_1}, \ldots, K_{a_k}$, $a_0 + a_1 + \cdots + a_k = n$. Without loss of generality, assume that $a_0 \geq a_1 \geq a_2 \geq \cdots \geq a_k \geq 1$.

If $a_0 = 1$, then G is an n-vertex tree. By [13, Theorems 3.3 and 3.8], we know the second last tree in an S-order among n-vertex trees is just $K_{1,n-1}^*$. It is easy to see that $a_0 \neq 2$, hence in what follows we consider $a_0 \geq 3$.

Let $V_i = \{v \in V_{K_{a_i}} : v \text{ is an end-vertex of a cut edge of } G\}$. In order to complete the proof, it suffices to show the following facts.

Fact 1. If $a_0 \geq 3$, then $|V_0| = 2, |V_1| = |V_2| = \cdots = |V_k| = 1$.

Proof. We prove Fact 1 by contradiction. If $|V_0| = |V_1| = |V_2| = \cdots = |V_k| = 1$, then without loss of generality assume that $V_i = \{u_i\},\ i = 0, 1, \ldots, k$.

First we consider $G \in \mathscr{K}_n^k$. Note that $G \in \mathscr{K}_n^k \setminus \{K_n^k\}$, hence $a_1 \geq 3$; otherwise, $a_1 = 2$, which implies that G contains at least $k+1$ cut edges, a contradiction. If $a_1 > 3$, we consider graph $G^* := K(a_0+1, \{a_1-1, a_2, \ldots, a_k\})$ in $\mathscr{K}_n^k \setminus \{K_n^k\}$. Note that $S_i(G) = S_i(G^*)$ for $i = 0, 1$ and $S_2(G) - S_2(G^*) = 2(a_1 - 1 - a_0) < 0$, hence $G \prec_s G^*$, a contradiction. Therefore $a_1 = 3$.

If $a_2 > 1$, we consider graph $G' := K(a_0 + a_2 - 1, \{3, 1, a_3, \ldots, a_k\}) \in \mathscr{G}_n^k \backslash K_n^k$. Note that $S_i(G) = S_i(G')$ for $i = 0, 1$ and $S_2(G) - S_2(G') = 2(a_2 - 1 - (a_2 - 1)a_0) = 2(a_2 - 1)(1 - a_0) < 0$, hence $G \prec_s G'$, a contradiction. Therefore, $a_2 = 1$, whence $a_3 = \cdots = a_k = 1$. Together with $a_1 = 3$, we have $a_0 = n-k-2$. That is to say, $G \cong K(n-k-2, \{3, 1, 1, \ldots, 1\})$.

For convenience, let $w_1 \in N_{K_{a_0}}(u_0)$ and $N_G(u_1) = \{u_0, v_1, v_2\}$. Consider

$$G^* := G - \{u_1v_1, u_1v_2\} + \{w_1v_1, w_1v_2\},$$

it is easy to see that $G^* \in \mathscr{G}_n^k \backslash K_n^k$. Note that $S_i(G) = S_i(G^*)$ for $i = 0, 1, 2, 3$ and $\phi_G(P_2) = \phi_{G^*}(P_2), \phi_G(C_4) = \phi_{G^*}(C_4)$, $\phi_G(P_3) - \phi_{G^*}(P_3) = 2 - 2(a_0 - 1) = 2(2 - a_0) < 0$, hence by Lemma 1.2(i) $S_4(G) < S_4(G^*)$. Thus $G \prec_s G^*$, a contradiction. Therefore $G \notin \mathscr{K}_n^k$.

Now we consider the case $G \notin \mathscr{K}_n^k$. It is easy to see that the edge induced graph $G[\mathbb{E}]$ is a tree which is not isomorphic to $K_{1,k}$. Hence, partition $V_{G[\mathbb{E}]}$ into $D^0(G[\mathbb{E}]) \cup D^1(G[\mathbb{E}]) \cup D^2(G[\mathbb{E}]) \cup D^3(G[\mathbb{E}]) \cup \cdots$, where $D^i(G[\mathbb{E}]) = \{u \in V_{G[\mathbb{E}]} : \text{dist}_{G[\mathbb{E}]}(u, u_0) = i\}$, $i = 0, 1, 2, 3, \ldots$. It is easy to see that $D^2(G[\mathbb{E}]) \neq \emptyset$.

If $D^3(G[\mathbb{E}]) \neq \emptyset$, that is to say, there exists $u \in D^2(G[\mathbb{E}])$ such that $d_{G[\mathbb{E}]}(u) \geq 2$, then choose u_i from $D^1(G[\mathbb{E}])$ such that u_i is adjacent to u_0 and u. Let $W := N_{G[\mathbb{E}]}(u_i) \setminus \{u_0\}$. As $u \in W$, we have $W \neq \emptyset$. Consider

$$G^* = G - \{u_i w : w \in W\} + \{u_0 w : w \in W\},$$

then its routine to check that $G^* \in \mathscr{G}_n^k \setminus \{K_n^k\}$. Note that $S_i(G) = S_i(G^*)$ for $i = 0, 1, 2, 3$ and $\phi_G(P_2) = \phi_{G^*}(P_2), \phi_G(C_4) = \phi_{G^*}(C_4)$, hence by Lemma 1.2(i) we have

$$S_4(G) - S_4(G^*) = 4(\phi_G(P_3) - \phi_{G^*}(P_3)) = 4[(a_i - a_0) - st],$$

where $s = |W| \geq 1$ and $t = |N_{G[\mathbb{E}]}(u_0) \setminus \{u_i\}| \geq 0$. Note that $a_i \leq a_0$, hence if $a_i < a_0$, then for all $t \geq 0$ we have $(a_i - a_0) - st < 0$, which implies that $G \prec_s G^*$, a contradiction. If $a_i = a_0$, then for all $t \geq 1$ we have $(a_i - a_0) - st < 0$, which implies that $G \prec_s G^*$, a contradiction. If $a_i = a_0$ and $t = 0$, then $G^* \cong G$. Hence, in order to complete the proof, it suffices to consider $D^3(G[\mathbb{E}]) = \emptyset$ and $d_{G[\mathbb{E}]}(u_0) > 1$. Furthermore, as $G \notin \mathscr{K}_n^k$, we have $D^2(G[\mathbb{E}]) \neq \emptyset$ and for each $u \in D^2(G[\mathbb{E}])$, u is a leaf of $G[\mathbb{E}]$ (otherwise, $D^3(G[\mathbb{E}]) \neq \emptyset$, a contradiction).

If there exists $u_i \in D^1(G[\mathbb{E}])$ such that $d_{G[\mathbb{E}]}(u_i) \geq 3$, then move $d_{G[\mathbb{E}]}(u_i) - 2$ pendant edges to u_0 and denote the resultant graph by G'. It is easy to see that $G' \in \mathscr{G}_n^k \setminus \{K_n^k\}$. Note that $s := d_{G[\mathbb{E}]}(u_i) - 2 \geq 1$, $q := d_{G[\mathbb{E}]}(u_0) - 1 \geq 1$, $S_i(G) = S_i(G')$ for $i = 0, 1, 2, 3$ and $\phi_G(P_2) = \phi_{G'}(P_2), \phi_G(C_4) = \phi_{G'}(C_4)$, hence by Lemma 1.2(i) we have

$$\begin{aligned} S_4(G) - S_4(G') &= 4(\phi_G(P_3) - \phi_{G'}(P_3)) \\ &= 4[s(a_i - 1) - s(a_0 - 1) - s(q - 1)] \\ &= 4s(a_i - a_0 - q + 1). \end{aligned} \tag{2.1}$$

If $a_0 > a_i$ or $q \geq 2$, in the view of (2.1), we obtain that $S_4(G) - S_4(G') < 0$, i.e., $G \prec_s G'$, a contradiction. If $a_0 = a_i$ and $q = 1$, then it is easy to see $G' \cong G$. Hence, in order to complete the proof, it suffices to consider that, in the edge induced graph $G[\mathbb{E}]$, each of the non-pendant vertices in $D^1(G[\mathbb{E}])$ is of degree 2.

For convenience, let $W = \{u : u \in D^1(G[\mathbb{E}]), d_{G[\mathbb{E}]}(u) = 2\}$. It is easy to see that $W \neq \emptyset$. If $|W| \geq 2$, choose $u \in W$ such that its unique neighbor in $G[\mathbb{E}]$ is a leaf, say u'. Let

$$G^* = G - uu' + u_0 u',$$

then $G^* \in \mathscr{G}_n^k \setminus \{K_n^k\}$. Since $S_i(G) = S_i(G^*)$ for $i = 0, 1, 2, 3$ and $\phi_G(P_2) = \phi_{G^*}(P_2), \phi_G(C_4) = \phi_{G^*}(C_4)$,

$$\phi_G(P_3) - \phi_{G^*}(P_3) = (a_i - 1) - (a_0 - 1) - p = a_i - a_0 - p < 0,$$

we have $S_4(G) - S_4(G^*) < 0$, i.e., $G \prec_s G^*$, a contradiction. Hence, $|W| = 1$.

By a similar discussion as in the proof of Fact 3 in Theorem 2.1, we can obtain that $a_0 = n - k, a_1 = a_2 = \ldots = a_k = 1$. Note that $a_0 \geq 3$, hence $k < n - 1$. Assume that $W = \{u\}$ with $N_{G[\mathbb{E}]}(u) = \{u_0, u'\}$, where u' is a pendant vertex in $G[\mathbb{E}]$. Let $x \in N_{K_{a_0}}(u_0)$. Consider $G^* = G - \{uu'\} + \{xu'\}$, then

$G^* \in \mathscr{G}_n^k \setminus \{K_n^k\}$. Note that $S_i(G) = S_i(G^*)$ for $i = 0, 1, 2, 3$, $\phi_G(P_2) = \phi_{G^*}(P_2), \phi_G(C_4) = \phi_{G^*}(C_4)$ and $\phi_G(P_3) - \phi_{G^*}(P_3) = 1 - (a_0 - 1) = 2 - a_0 < 0$ ($a_0 \geq 3$), hence $S_4(G) - S_4(G^*) < 0$, i.e., $G \prec_s G^*$, a contradiction.

If there is a V_i satisfying $|V_i| \geq 3$, then choose two distinct vertices u_i', u_i'' in V_i and let $G^* = G - \{u_i'u : u \in N_{G[\mathbb{E}]}(u_i')\} + \{u_i''u : u \in N_{G[\mathbb{E}]}(u_i')\}$. It is easy to see that $G^* \in \mathscr{G}_n^k \setminus \{K_n^k\}$. Note that $S_i(G) = S_i(G^*), i = 0, 1, 2, 3,\quad \phi_G(P_2) = \phi_{G^*}(P_2), \phi_G(C_4) = \phi_{G^*}(C_4)$ and $\phi_G(P_3) - \phi_{G^*}(P_3) = -|N_{G[\mathbb{E}]}(u_i')||N_{G[\mathbb{E}]}(u_i'')| < 0$, hence $S_4(G) - S_4(G^*) < 0$, which implies that $G \prec_s G^*$, a contradiction.

If there exists an $i \in \{1, 2, \ldots, k\}$ such that $|V_i| = 2$. Assume, without loss of generality, that $V_i = \{u_i, u'\}$, where u_i is adjacent to $u_0 \in V_0$. Let

$$G^* = G - \{u_i x : x \in V_{K_{a_i}}\} + \{yx : y \in V_{K_{a_0}}, x \in V_{K_{a_i}} \setminus \{u_i\}\}.$$

It is easy to see that $G^* \in \mathscr{G}_n^k \backslash K_n^k$. Note that $S_i(G) = S_i(G^*)$ for $i = 0, 1$ and $S_2(G) - S_2(G^*) = 2(a_i - 1) - 2(a_i - 1)a_0 = 2(a_i - 1)(1 - a_0) < 0$, hence $S_2(G) < S_2(G^*)$, i.e., $G \prec_s G^*$, a contradiction.

Combining with discussion as above, we obtain that $|V_1| = |V_2| = \cdots = |V_k| = 1$, whence $|V_0| = 2$, as desired. □

Fact 2. $a_1 = a_2 = \cdots = a_k = 1$.

Proof. By a similar discussion as in the proof of Fact 3 in Theorem 2.1, we can get $a_0 = n - k$, $a_1 = a_2 = \cdots = a_k = 1$. We omit the procedure here. □

Fact 3. $G \cong G_3$, where $G_3 = K(n - k, \{1, 1, \ldots, 1\}) - u_0u_k + u_0'u_k$, where u_0u_k is a cut edge and $u_0' \in V_{K_{a_0}} \backslash \{u_0\}$.

Proof. Note that if G has just two cut edges, it is easy to see that $G \cong G_3$ defined as above. Hence in what follows we consider that G contains at least three cut edges.

Let $N_{G[\mathbb{E}]}(u_0) = \{u_1, u_2, \ldots, u_m\}$ and $N_{G[\mathbb{E}]}(u_0') = \{u_1', u_2', \ldots, u_t'\}$. Without loss of generality, assume that $m \geq t$. Obviously, $t \geq 1$. At first we show that $t = 1$. Otherwise, let

$$G^* = G - \{u_0'u_2', u_0'u_3', \ldots, u_0'u_t'\} + \{u_0u_2', u_0u_3', \ldots, u_0u_t'\}.$$

It is easy to see that $G^* \in \mathscr{G}_n^k \backslash K_n^k$. Note that $S_i(G) = S_i(G^*)$ for $i = 0, 1, 2, 3$, $\phi_G(P_2) = \phi_{G^*}(P_2)$, and $\phi_G(C_4) = \phi_{G^*}(C_4)$, hence

$$S_4(G) - S_4(G^*) = 4(\phi_G(P_3) - \phi_{G^*}(P_3)) = -4m(t - 1) < 0,$$

i.e., $G \prec_s G^*$, a contradiction.

Now we are to show that $m = k - 1$. If not, there exists a vertex $u \in \{u_1, u_2, \ldots, u_m, u_1', u_2', \ldots, u_t'\}$ such that $d_{G[\mathbb{E}]}(u) \geq 2$. Denote $N_{G[\mathbb{E}]}(u) \setminus \{u_0, v_0\} = \{\hat{u}_1, \hat{u}_2, \ldots, \hat{u}_s\}$, $s \geq 1$. Let

$$G^* = G - \{u\hat{u}_1, u\hat{u}_2, \ldots, u\hat{u}_s\} + \{u_0\hat{u}_1, u_0\hat{u}_2, \ldots, u_0\hat{u}_s\}.$$

It is easy to see that $G^* \in \mathscr{G}_n^k \backslash K_n^k$. Notice that $S_i(G) = S_i(G^*)$ for $i = 0, 1, 2, 3$, $\phi_G(P_2) = \phi_{G^*}(P_2)$ and $\phi_G(C_4) = \phi_{G^*}(C_4)$, hence

$$S_4(G) - S_4(G^*) = 4(\phi_G(P_3) - \phi_{G^*}(P_3)) = 4s((a_i - 1) - (a_0 - 1) - (m - 1)) = 4s(a_i - a_0) - 4s(m - 1) < 0.$$

The last inequality follows by $a_i = 1 < n - k = a_0$ (by fact 2), and $m \geq 1, s \geq 1$. Hence, we get $S_4(G) < S_4(G^*)$, i.e., $G \prec_s G^*$, a contradiction. so we have $m = k - 1, t = 1$, which is equivalent to that $G \cong G_3$. □

This completes the proof. $\square$

3. The first and the second graphs in an S-order among $\mathscr{G}_n^k$

In this section, we are to determine the first and the second graphs in an S-order among $\mathscr{G}_n^k$. Let $\mathbb{E} = \{e_1, e_2, \dots, e_k\}$ be the set of all the cut edges of $G \in \mathscr{G}_n^k$. Note that if we delete an edge, say e, from a connected graph G, then in the view of $S_2(G) = 2|E_G|$, we have $S_2(G) > S_2(G - e)$. In order to determine the first graph in an S-order among $\mathscr{G}_n^k$, it suffices to choose the graph such that its size is as small as possible.

Theorem 3.1. *Of all the connected graphs with n vertices and k cut edges, the first graph in an S-order is obtained uniquely at P_n^k.*

Proof. Choose $G \in \mathscr{G}_n^k$ such that it is as small as possible according to the relation $\preceq_s$. If $k = 0$, then it is easy to see that $G \cong C_n$ and our result holds immediately. Therefore we may assume that $k \geq 1$. We show the following claim at first.

Claim 1. *G contains exactly one cycle.*

Proof. Assume to the contrary that G contains at least two cycles. If G contains two cycles C^1 and C^2 such that C^1 and C^2 have edges in common; see Fig. 2(a), then let $G^* = G - \{uv, xy\} + ux$ (see Fig. 2(b)); if G contains two cycles C^1 and C^2 such that C^1 and C^2 have just one vertex in common; see Fig. 2(c), then let $G^* = G - \{ux, vx\} + uv$ (see Fig. 2(d)). It is routine to check that $G^* \in \mathscr{G}_n^k$ and in each of the above cases one has $S_i(G) = S_i(G^*)$, $i = 0, 1$ and $S_2(G) - S_2(G^*) = 2 > 0$, hence $G^* \prec_s G$, a contradiction.



Figure 2: Graphs used in the proof of Claim 1.

If G contains two cycles $C_l = u_0u_1u_2 \dots u_{l-1}$ and $C_j = v_0v_1v_2 \dots v_{j-1}$ such that C_l connects C_j by a path P_i, $i \geq 2$, whose end vertices are u_0, v_1, and the vertex, say u_t (resp. v_m), on the cycle C_l (resp. C_j) in G either is of degree 2 or has subgraph G_t (resp. H_m) attached, $0 \leq t \leq l-1$, $0 \leq m \leq j-1$; see Fig. 3. Let

$$G^* = G - \{u_0u_1, v_1v_2, v_0v_1\} + \{u_0v_2, u_1v_0\},$$

then $G^* \in \mathscr{G}_n^k$. Since $S_i(G) = S_i(G^*)$, $i = 0, 1$. $S_2(G) - S_2(G^*) = 2 > 0$, then $G^* \prec_s G$, a contradiction. Therefore, G contains exactly one cycle. $\square$

By Claim 1, we know that G is a unicyclic graph. Note that G contains exactly k cut edges, hence G is an n-vertex unicyclic graph with girth $n - k$. By Lemma 1.4 the first graph in an S-order among the n-vertex unicyclic graph with girth $n - k$ is just the graph P_n^k, as desired. $\square$

At the rest of this section, we are to determine the second graph in an S-order among $\mathscr{G}_n^k$ $(k \geq 3)$.



Figure 3: Graph $G \Rightarrow G^*$.

Theorem 3.2. *Of all graphs with* n *vertices and* k *cut edges, the second graph in an* S*-order is obtained uniquely at* $\hat{U}_n^k$ $(k \geq 3)$*, where* $\hat{U}_n^k$ *is obtained by attaching two leafs to the pendant vertex of graph* P_{n-2}^{k-2}*.*

Proof. Note that if we delete an edge e from a connected graph G, then in the view of $S_2(G) = 2|E_G|$, we have $S_2(G) > S_2(G-e)$, hence in order to determine the second graph in an S-order among $\mathscr{G}_n^k$, it suffices to determine the second graph in an S-order among the set of all n-vertex unicyclic graphs with girth $n-k$; we denote this set by $\mathscr{U}_n^k$.

Choose $G \in \mathscr{U}_n^k \setminus \{P_n^k\}$ such that it is as small as possible with respect to $\preceq_s$. Note that $\mathbb{E}$ is the set of k cut edges of G, hence $G[\mathbb{E}]$ is a forest. We are to show that $G[\mathbb{E}]$ is a tree. If this is not true, then it is equivalent to that there exist at least two vertices, say u_0, v_0, on the unique cycle contained in G satisfying $d_G(u_0), d_G(v_0) \geq 3$.

In the edge induced graph $G[\mathbb{E}]$, consider the tree, say T_1, containing u_0. We are to show that T_1 is a path; otherwise, choose a longest path $P = u_0u_1 \ldots u_p$ in T_1 with end-vertex u_0, u_p, it is easy to see $d_{T_1}(u_p) = 1$. If there exists u_i with $i \geq 1$ on P such that $d_G(u_i) > 2$. Choose a vertex x in $N_G(u_i) \setminus \{u_{i-1}, u_{i+1}\}$ and let $G^* = G - u_ix + u_px$, then $G^* \in \mathscr{G}_n^k \backslash \{P_n^k\}$. Note that $S_i(G) = S_i(G^*)$ for $i = 0, 1, 2, 3$, $\phi_G(P_2) = \phi_{G^*}(P_2), \phi_G(C_4) = \phi_{G^*}(C_4)$ and $\phi_G(P_3) - \phi_{G^*}(P_3) \geq 1$, hence by Lemma 1.2(i), we get $S_4(G) - S_4(G^*) > 0$, i.e., $G^* \prec_s G$, a contradiction. Hence, we obtain that each vertex u_i on P is of degree 2 in G for $i = 1, 2, \ldots, p-1$. Hence, if $d_G(u_0) = 3$, then T_1 is a path, as desired. If $d_G(u_0) > 3$, then choose x from $N_G(u_0)$ such that x is not on the cycle and the path P contained in G. Let $G^* = G - u_0x + u_px$, then $G^* \in \mathscr{G}_n^k \backslash P_n^k$. Notice that $S_i(G) = S_i(G^*)$ for $i = 0, 1, 2, 3$, $\phi_G(P_2) = \phi_{G^*}(P_2), \phi_G(C_4) = \phi_{G^*}(C_4)$ and $\phi_G(P_3) - \phi_{G^*}(P_3) \geq 2$, by Lemma 1.2(i), we get $S_4(G) - S_4(G^*) > 0$, i,e., $G^* \prec_s G$, a contradiction. By a similar discussion as above, we can also show that, in $G[\mathbb{E}]$, the component contains v_0 is also a path, say P'. For convenience, let v_0' be the neighbor of v_0 on P'.

If there exists another vertex $u_0' \neq u_0, v_0$, on the unique cycle contained in G satisfying $d_G(u_0') \geq 3$. Let $G^* = G - \{u_0'x, x \in N_{G[\mathbb{E}]}(u_0')\} + \{u_px, x \in N_{G[\mathbb{E}]}(u_0')\}$, then $G^* \in \mathscr{G}_n^k \backslash P_n^k$. Note that $S_i(G) = S_i(G^*)$ for $i = 0, 1, 2, 3$, $\phi_G(P_2) = \phi_{G^*}(P_2), \phi_G(C_4) = \phi_{G^*}(C_4)$ and $\phi_G(P_3) - \phi_{G^*}(P_3) \geq 1$, by Lemma 1.2(i), we get $S_4(G) - S_4(G^*) > 0$, i,e., $G^* \prec_s G$, a contradiction. So, we just need to consider there exist two vertices on the unique cycle contained in G satisfying $d_G(u_0), d_G(v_0) \geq 3$. Without loss of generality, assume that $|E_P| \geq |E_{P'}|$. Let $G^* = G - v_0v_0' + u_{p-1}v_0'$, it is easy to see that $G^* \in \mathscr{G}_n^k \setminus \{P_n^k\}$.

- $k = 3$. By Lemma 1.1, we have $S_i(G) \geq S_i(G^*)$ for $i = 0, 1, \ldots, n-2$ and $S_{n-1}(G) > S_{n-1}(G^*)$. Hence $G^* \prec_s G$, a contradiction.
- $k \geq 4$. Note that $\phi_G(P_2) = \phi_{G^*}(P_2), \phi_G(C_4) = \phi_{G^*}(C_4)$, $\phi_G(P_3) = \phi_{G^*}(P_3), \phi_G(K_{1,3}) = \phi_{G^*}(K_{1,3})$, $\phi_G(U_5) = \phi_{G^*}(U_5), \phi_G(B_4) = \phi_{G^*}(B_4)$, $\phi_G(B_5) = \phi_{G^*}(B_5)$, $\phi_G(C_3) = \phi_{G^*}(C_3)$, $\phi_G(C_6) = \phi_{G^*}(C_6)$ and $\phi_G(P_4) - \phi_{G^*}(P_4) \geq 1$, hence by Lemma 1.2, 1.3, we get that $S_i(G) = S_i(G^*)$ for $i = 0, 1, 2, 3, 4, 5$ and $S_6(G) - S_6(G^*) > 0$, i.e., $G^* \prec_s G$, a contradiction.

Therefore, we obtain that $G[\mathbb{E}]$ is a tree. That is to say, there exists just one vertex, say u_0, on the unique cycle such that $d_G(u_0) \geq 3$. Choose one of the longest paths, say $P := u_0u_1 \ldots u_p$, from $G[\mathbb{E}]$. It is easy to see that u_p is a leaf of G. Furthermore, we have the following claim.

Claim 2. *The length of P is $k-1$, i.e., $P := u_0u_1 \dots u_{k-2}u_{k-1}$ and $G[\mathbb{E}]$ is obtained from P by attaching a leaf to u_{k-2} of P.*

Proof. Note that $P = u_0u_1 \dots u_p$ is one of the longest paths of $G[\mathbb{E}]$ and u_p is a leaf. Hence, we first show that $d_G(u_0) = 3$. Otherwise, choose x from $N_G(u_0)$ such that x is not on the cycle and the path P of G. If $d_G(u_i) \geq 3$ for some $i \in \{1, 2, \dots, p-1\}$, let $G^* = G - u_0x + u_px$. Obviously, $G^* \in \mathscr{G}_n^k \backslash \{P_n^k\}$. Note that $S_i(G) = S_i(G^*)$, $i = 0, 1, 2, 3$, $\phi_G(P_2) = \phi_{G^*}(P_2), \phi_G(C_4) = \phi_{G^*}(C_4)$ and $\phi_G(P_3) - \phi_{G^*}(P_3) \geq 2$, hence by Lemma 1.2(i), we get $S_4(G) - S_4(G^*) > 0$, i.e., $G^* \prec_s G$, a contradiction. If $d_G(u_i) = 2$ for any $i \in \{1, 2, \dots, p-1\}$, let $G^* = G - u_0x + u_{p-1}x$. Obviously, $G^* \in \mathscr{G}_n^k \backslash \{P_n^k\}$. Note that $S_i(G) = S_i(G^*)$, $i = 0, 1, 2, 3$, $\phi_G(P_2) = \phi_{G^*}(P_2), \phi_G(C_4) = \phi_{G^*}(C_4)$ and $\phi_G(P_3) - \phi_{G^*}(P_3) \geq 1$, hence by Lemma 1.2(i), we get $S_4(G) - S_4(G^*) > 0$, i.e., $G^* \prec_s G$, a contradiction.

Now we show that $d_G(u_i) = 2$, $i = 1, 2, \dots, p-2$ and $d_G(u_{p-1}) = 3$. Note that $G \ncong P_n^k$, hence there exists at least one vertex u_i $(1 \leq i \leq p-1)$ on P such that $d_G(u_i) \geq 3$.

If there exists a vertex u_i $(1 \leq i \leq p-1)$ on P such that $d_G(u_i) \geq 4$, then choose $x \in N_G(u_i) \setminus \{u_{i-1}, u_{i+1}\}$ and let $G^* = G - u_ix + u_px$. Obviously, $G^* \in \mathscr{G}_n^k \backslash P_n^k$. Since $S_i(G) = S_i(G^*)$, $i = 0, 1, 2, 3$. $\phi_G(P_2) = \phi_{G^*}(P_2)$, $\phi_G(C_4) = \phi_{G^*}(C_4)$ and $\phi_G(P_3) - \phi_{G^*}(P_3) \geq 2$, by Lemma 1.2(i), we have $S_4(G) - S_4(G^*) > 0$, i.e., $G^* \prec_s G$, a contradiction. Hence, $\max\{d_G(u_i), i = 1, 2, \dots, p-1\} = 3$.

If $d_G(u_{p-1}) = d_G(u_i) = 3$ for some $i \in \{1, 2, \dots, p-2\}$, then choose z_1 in $N_G(u_i) \setminus \{u_{i-1}, u_{i+1}\}$ and let $G^* = G - u_iz_1 + u_pz_1$, then $G^* \in \mathscr{G}_n^k \backslash P_n^k$. Notice that $S_i(G) = S_i(G^*)$, $i = 0, 1, 2, 3$. $\phi_G(P_2) = \phi_{G^*}(P_2)$, $\phi_G(C_4) = \phi_{G^*}(C_4)$ and $\phi_G(P_3) - \phi_{G^*}(P_3) = 1 > 0$, hence by Lemma 1.2(i), we get that $S_4(G) - S_4(G^*) > 0$, i.e., $G^* \prec_s G$, a contradiction.

If $d_G(u_{p-1}) = 2$, $d_G(u_i) = 3$ for some $i \in \{1, 2, \dots, p-2\}$, then choose z_1 in $N_G(u_i) \setminus \{u_{i-1}, u_{i+1}\}$ and let $G^* = G - u_iz_1 + u_{p-1}z_1$, then it is easy to see that $G^* \in \mathscr{G}_n^k \backslash \{P_n^k\}$. Note that $S_i(G) = S_i(G^*)$ for $i = 0, 1, 2, 3, 4, 5$, $\phi_G(P_2) = \phi_{G^*}(P_2)$, $\phi_G(C_4) = \phi_{G^*}(C_4)$, $\phi_G(P_3) = \phi_{G^*}(P_3)$, $\phi_G(K_{1,3}) = \phi_{G^*}(K_{1,3})$, $\phi_G(U_5) = \phi_{G^*}(U_5)$, $\phi_G(B_4) = \phi_{G^*}(B_4)$, $\phi_G(B_5) = \phi_{G^*}(B_5)$, $\phi_G(C_3) = \phi_{G^*}(C_3)$, $\phi_G(C_6) = \phi_{G^*}(C_6)$ and $\phi_G(P_4) - \phi_{G^*}(P_4) \geq 1$, hence by Lemma 1.2(iii), we get that $S_6(G) - S_6(G^*) > 0$, i.e., $G^* \prec_s G$, a contradiction.

Hence, we obtain that $d_G(u_0) = 3, d_G(u_1) = d_G(u_2) = \dots = d_G(u_{p-2}) = 2, d_G(u_{p-1}) = 3$ and $d_G(u_p) = 1$. For convenience, let $N_G(u_{p-1}) \setminus \{u_{p-2}, u_p\} = \{z_0\}$. It is easy to see that z_0 is a leaf; otherwise $G[\mathbb{E}]$ contains a path $P' := u_0u_1 \dots u_{p-1}z_0 \dots z_t$, where z_t is a leaf. It is routine to check that the length of P' is longer than that of P, a contradiction. Therefore, $d_G(z_0) = 1$, as desired. □

Based on Claim 2, Theorem 3.2 follows immediately. □

References

[1] B.A. Bondy, U.S.R. Murty, Graph Theory with Applications, Macmillan, London, 1976.

[2] B. Cheng, B.L. Liu, J.X. Liu, On the spectral moments of unicyclic graphs with fixed diameter. Linear Algebra Appl. 437 (4) (2012) 1123-1131.

[3] D. Cvetković, M. Doob, H. Sachs, Spectral of Graphs-Theory and Applications, Academic Press, New York, 1980.

[4] D. Cvetković, M. Doob, H. Sachs, A. Torgaśev, Recent Results in the Theory of Graph Spectral, Annals of Discrete Mathematics Series, North-Holland, 1988.

[5] D. Cvetković, M. Petrić, A table of connected graphs on six vertices, Discrete Math. 50 (1984) 37-49.

[6] D. Cvetković, P. Rowlinson, Spectra of unicyclic graphs, Graphs Comb. 3 (1987) 7-23.

[7] Y.Q. Feng, X. Hu, S.C. Li, On the extremal Zagreb indices of graphs with cut edges, Acta Appl. Math. 110 (2010) 667-684.

[8] S.N. Hu, S.C. Li, X.X. Zhang, On the spectral moment of graphs with given clique number, submission.

[9] S.C. Li, Y.B. Song, On the spectral moment of trees with given degree sequences, arXiv:1209.2188v1 [math.CO] 11 Sep 2012.

[10] X.F. Pan, X.G. Liu, H.Q. Liu, On the spectral moment of quasi-trees, Linear Algebra Appl. 436 (2012) 927-934.

[11] X.F. Pan, X.L. Hu, X.G. Liu, H.Q. Liu, The spectral moments of trees with given maximum degree, Appl. Math. Lett. 24 (7) (2011) 1265-1268.

[12] Y.P. Wu, Q. Fan, On the lexicographical ordering by spectral moments of bicyclic graphs, Ars Combin., in press.

[13] Y.P. Wu, H.Q. Liu, Lexicographical ordering by spectral moments of trees with a prescribed diameter, Linear Algebra Appl. 433 (2010) 1707-1713.

[14] H.Liu, M. Lu, F. Tian, On the spectral radius of graphs with cut edges. Linear Algebra Appl. 389, 139-145 (2004)

[15] R. Pepper, G. Henry, D. Sexton, Cut-edges and the independence number. MATCH Commun. Math. Comput. Chem. 56, 403-408 (2006)

[16] Y.R. Wu, S. He, J.L. Shu, Largest spectral radius among graphs with k cut edges. J. East China Norm. Univ. Nat. Sci. Ed. 3, 67-74 (2007) (Chinese)